

A 2–Cocycle on a Group of Symplectomorphisms

Rais S. Ismagilov
Mark Losik
Peter W. Michor

Vienna, Preprint ESI 1525 (2004)

February 24, 2006

Supported by the Austrian Federal Ministry of Education, Science and Culture
Available via <http://www.esi.ac.at>

A 2-COCYCLE ON A GROUP OF SYMPLECTOMORPHISMS

RAIS S. ISMAGILOV, MARK LOSIK, PETER W. MICHOR

ABSTRACT. For a symplectic manifold (M, ω) with exact symplectic form we construct a 2-cocycle on the group of symplectomorphisms and indicate cases when this cocycle is not trivial.

1. INTRODUCTION

For a symplectic manifold (M, ω) such that $H^1(M, \mathbb{R}) = 0$ and the symplectic form ω is exact we indicate a formula defining a 2-cocycle on the group $\text{Diff}(M, \omega)$ of symplectomorphisms with values in the trivial $\text{Diff}(M, \omega)$ -module \mathbb{R} . Let G be a connected real simple Lie group and K a maximal compact subgroup. For the symmetric Hermitian space $M = G/K$ endowed with the induced symplectic structure, we prove that the restriction of this cocycle to the group G is non-trivial. Thus this cocycle is non-trivial on the whole group $\text{Diff}(M, \omega)$, too. In particular, this implies that the cocycle is non-trivial for the symplectic manifold $(\mathbb{R}^2 \times M, \omega_0 + \omega_M)$, where (M, ω_M) is a non-compact symplectic manifold with exact symplectic form ω_M such that $H^1(M, \mathbb{R}) = 0$ and ω_0 is the standard symplectic form on \mathbb{R}^2 .

For the convenience of the reader, in an appendix we consider the corresponding 2-cocycle on the Lie algebra of locally Hamiltonian and Hamiltonian vector fields and indicate when this cocycle is non-trivial.

Note that in [7] a similar 2-cocycle was constructed for the group of volume preserving diffeomorphisms on a compact n -dimensional manifold M . This cocycle takes its values in the space $H^{n-2}(M, \mathbb{R})$. Neretin in [10] constructed a 2-cocycle on the group of symplectomorphisms with compact supports.

Throughout the paper M is a connected C^∞ -manifold.

We thank Stefan Haller and Yurii Neretin for comments.

Date: February 24, 2006.

2000 Mathematics Subject Classification. Primary 58D05, 20J06, 22E65.

Key words and phrases. group extension, symplectomorphism.

ML and PWM were supported by FWF Projekt P 17108-N04. PWM was supported by Centre Bernoulli, Lausanne.

2. PRELIMINARIES

We recall some standard facts on central extensions of groups and two-dimensional cohomology of groups (see, for example, [8], ch. 4).

Consider a group G and the field \mathbb{R} as a trivial G -module. Let $C^p(G, \mathbb{R})$ be the set of maps from G^p to \mathbb{R} for $p > 0$ and let $C^0(G, \mathbb{R}) = \mathbb{R}$. Define a map $D^p : C^p(G, \mathbb{R}) \rightarrow C^{p+1}(G, \mathbb{R})$ as follows: for $f \in C^p(G, \mathbb{R})$ and $g_1, \dots, g_{p+1} \in G$

$$(1) \quad (D^p f)(g_1, \dots, g_{p+1}) = f(g_2, \dots, g_{p+1}) \\ + \sum_{i=1}^p (-1)^i f(g_1, \dots, g_i g_{i+1}, \dots, g_{p+1}) + (-1)^{p+1} f(g_1, \dots, g_p).$$

By definition $C^*(G, \mathbb{R}) = (C^p(G, \mathbb{R}), D^p)_{p \geq 0}$ is the standard complex of nonhomogeneous cochains of the group G with values in the G -module \mathbb{R} and its cohomology $H^*(G, \mathbb{R}) = (H^p(G, \mathbb{R}))_{p \geq 0}$ is the cohomology of the group G with values in the trivial G -module \mathbb{R} . Recall that a cochain $f \in C^p(G, \mathbb{R})$ is called normalized if $f(g_1, \dots, g_p) = 0$ whenever at least one of the $g_1, \dots, g_p \in G$ equals the identity e of G . It is known that the inclusion of the subcomplex of normalized cochains into $C^*(G, \mathbb{R})$ induces an isomorphism in cohomology.

Let f be a normalized 2-cocycle of G with values in a trivial G -module \mathbb{R} . Let $E(G, \mathbb{R}) = G \times \mathbb{R}$, with multiplication $(g_1, a_1)(g_2, a_2) = (g_1 g_2, a_1 g_2 + a_2 + f(g_1, g_2))$ for $a_1, a_2 \in \mathbb{R}$ and $g_1, g_2 \in G$. Then $E(G, \mathbb{R})$ is a group, and the natural projection $E(G, \mathbb{R}) = G \times \mathbb{R} \rightarrow G$ is a central extension of the group G by \mathbb{R} . The extension $E(G, \mathbb{R})$ is non-split iff the cocycle f is non-trivial.

If G is a topological group (finite-dimensional or infinite-dimensional Lie group) one can define a subcomplex $C_{\text{cont}}^*(G, \mathbb{R})$ ($C_{\text{diff}}^*(G, \mathbb{R})$) of the complex $C^*(G, \mathbb{R})$ (see [4], ch. 3) consisting of cochains which are continuous (smooth) functions. The cohomologies of the complexes $C_{\text{cont}}^*(G, \mathbb{R})$ and $C_{\text{diff}}^*(G, \mathbb{R})$ are isomorphic whenever G is a finite-dimensional Lie group (see [4], ch. 3 and [9]). Note that if the 2-cocycle f is continuous (differentiable), the extension $E(G, \mathbb{R})$ is a topological group (Lie group).

3. A 2-COCYCLE ON THE GROUP OF SYMPLECTOMORPHISMS

Let (M, ω) be a non-compact symplectic manifold such that $H^1(M, \mathbb{R}) = 0$ and the symplectic form ω is exact. Let ω_1 be a 1-form on M such that $d\omega_1 = \omega$. Denote by $\text{Diff}(M, \omega)$ the group of symplectomorphisms of M . We define a 2-cocycle on the group $G = \text{Diff}(M, \omega)$ with values in the trivial G -module \mathbb{R} as follows. Fix a point $x_0 \in M$. Then for $g_1, g_2 \in G$ we put

$$(2) \quad C_{x_0}(g_1, g_2) = \int_{x_0}^{g_2 x_0} (g_1^* \omega_1 - \omega_1),$$

where the integral is taken along a smooth curve connecting the point x_0 with the point g_2x_0 . Since $H^1(M, \mathbb{R}) = 0$ the 1-form $g_1^*\omega_1 - \omega_1$ is exact and the value of this integral does not depend on the choice of such a curve.

Theorem 3.1. *The function $C_{x_0} : G^2 \rightarrow \mathbb{R}$ defined by (2) is a normalized 2-cocycle on the group G with values in the trivial G -module \mathbb{R} . The cohomology class of C_{x_0} is independent of the choice of the point x_0 and the form ω_1 .*

Proof. By (1) it is easy to check that $D^2C_{x_0} = 0$. Moreover, the 2-cocycle C_{x_0} is normalized. Since for each $g \in G$ the 1-form $g^*\omega_1 - \omega_1$ is exact, for any points $x_1, x_2 \in M$ we have $C_{x_1} - C_{x_2} = Da$, where a is a 1-cochain on G defined by $a(g) = \int_{x_1}^{x_2} (g^*\omega_1 - \omega_1)$. \square

By definition, the cocycle C_{x_0} is a continuous function on $G \times G$.

Remark 3.2. Let M be a manifold such that $H^1(M, \mathbb{R}) = 0$ and let ω be an exact 2-form on M . Let $\text{Diff}(M, \omega)$ be the group of diffeomorphisms of M preserving the form ω . Then the formula (2) for $g_1, g_2 \in \text{Diff}(M, \omega)$ gives a 2-cocycle on the group $\text{Diff}(M, \omega)$ and all statements of theorem (3.1) are true for this cocycle.

Denote by $E(\text{Diff}(M, \omega))$ the central extension of the group $\text{Diff}(M, \omega)$ by \mathbb{R} defined by the cocycle C_{x_0} . Now we give a geometric interpretation of the extension $E(\text{Diff}(M, \omega))$. We choose a form ω_1 with $d\omega_1 = \omega$ and put $\omega_2(g) = \int_{x_0}^x (\omega_1 - g^*\omega_1)$. Consider the trivial \mathbb{R} -bundle $M \times \mathbb{R}$. Clearly, the form $dt + \omega_1$ is a connection with curvature ω of this bundle. Denote by $\text{Aut}(M \times \mathbb{R}, \omega)$ the group of those bundle automorphisms which respect the connection $dt + \omega_1$ and which are projectable to diffeomorphisms in $\text{Diff}(M, \omega)$. It is easy to check that the group $\text{Aut}(M \times \mathbb{R}, \omega)$ is isomorphic to the group $E(\text{Diff}(M, \omega)) = \text{Diff}(M, \omega) \times \mathbb{R}$ which acts as follows on $M \times \mathbb{R}$: $(x, t) \rightarrow (g(x), \omega_2(g)(x) + t + a)$, where $(x, t) \in M \times \mathbb{R}$ and $(g, a) \in G \times \mathbb{R}$. This gives an equivalent definition of the extension $E(\text{Diff}(M, \omega))$ as a group of automorphisms of the trivial principal \mathbb{R} -bundle $M \times \mathbb{R}$ with connection $dt + \omega_1$.

If we replace the form ω_1 by the form $\omega_1 + df$, where f is a smooth function on M , we get an action of G on $M \times \mathbb{R}$ which is related to the initial one by the gauge transformation $(x, t) \rightarrow (x, t - f(x))$ of the bundle $M \times \mathbb{R} \rightarrow M$.

4. EXAMPLES OF NON-TRIVIAL 2-COCYCLES

The authors are not able to prove that the cocycle C_{x_0} is non-trivial for any symplectic manifold M with an exact symplectic 2-form ω . In this section we prove that for some symplectic manifolds the restrictions of this cocycle to some subgroups of $G \subset \text{Diff}(M, \omega)$ turn out to be non-trivial.

4.1. The linear symplectic space \mathbb{R}^{2n} and the Heisenberg group. Consider the space \mathbb{R}^{2n} with the standard symplectic form $\omega_0 = \sum_{k=1}^n dx_k \wedge dx_{k+n}$ and the group $G = \mathbb{R}^{2n}$ acting on the space \mathbb{R}^{2n} by translations. Applying (2) to the form

ω_0 , the 1-form $\omega_1 = \frac{1}{2} \sum_{k=1}^n (x_{n+k} dx_k - x_k dx_{n+k})$, and the point $x_0 = 0 \in \mathbb{R}^{2n}$ we get a 2-cocycle on the group G given by

$$C_0(x, y) = \frac{1}{2} \sum_{k=1}^n (x_k y_{n+k} - y_k x_{n+k}),$$

where $x = (x_1, \dots, x_{2n})$ and $y = (y_1, \dots, y_{2n})$. The central extension of the group \mathbb{R}^{2n} by \mathbb{R} defined by this cocycle is the Heisenberg group. This extension is non-split since the Heisenberg group is noncommutative and thus the cocycle $C_0(x, y)$ is non-trivial.

4.2. Symmetric Hermitian spaces and the Guichardet-Wigner cocycle.

Consider a non-compact symmetric space $M = G/K$, where G is a connected real simple Lie group and where K is a maximal compact subgroup. Then M is diffeomorphic to \mathbb{R}^n , where $n = \dim M$. We suppose that M admits a G -invariant complex structure, i.e., M is a symmetric Hermitian space. This condition is satisfied (up to finite covering) for the following groups: $SU(p, q)$ ($p, q \geq 1$), $SO_0(2, q)$ ($q = 1$ or $q \geq 3$), $Sp(n, \mathbb{R})$ ($n \geq 1$), $SO^*(2n)$ ($n \geq 2$), and certain real forms of E_6 and E_7 .

Consider the symplectic manifold (M, ω) , where the symplectic form ω is defined by the Hermitian metric on M . It is known that on each of the Lie groups mentioned above, in the complex $C_{\text{diff}}(G, \mathbb{R})$ there is a non-trivial Guichardet-Wigner 2-cocycle (see [5] and [4]). By [2] this cocycle is given as follows, up to a nonzero factor:

$$(3) \quad (g_1, g_2) \mapsto \int_{(x_0, g_1 x_0, g_1 g_2 x_0)} \omega,$$

where $g_1, g_2 \in G$, $x_0 = K \in G/K$, and the integral is taken over the oriented geodesic cone with vertex x_0 and the segment of a geodesic from $g_1 x_0$ to $g_1 g_2 x_0$ as base.

We prove that the restriction of the cocycle C_{x_0} to the group G is cohomologous to the cocycle given by (3).

For the base point x_0 we define a 1-cochain γ_{x_0} on the group G as follows:

$$\gamma_{x_0}(g) = \int_{x_0}^{g x_0} \omega_1,$$

where $g \in G$ and the integral is taken along the geodesic segment from x_0 to $g x_0$. Consider C_{x_0} on G given by formula (2), where we choose for the curve between the points x_0 and $g_2 x_0$ a geodesic segment from x_0 to $g_2 x_0$. It is easy to check that on the group G the cocycle $C_{x_0} + D\gamma_{x_0}$ equals the cocycle given by (3). Thus the cocycle C_{x_0} on the group G is non-trivial in the complex $C_{\text{diff}}^*(G, \mathbb{R})$.

In particular, for the group $G = SL(2, \mathbb{R}) = SU(1, 1)$ the symmetric space $M = G/K$ is the hyperbolic plane H^2 and ω is the area form on H^2 . Instead of the group $SL(2, \mathbb{R})$ we will later consider the group $PSL(2, \mathbb{R})$ which acts effectively on H^2 . Since $SL(2, \mathbb{R})$ is a two-sheet cover of $PSL(2, \mathbb{R})$, the cohomologies of these groups

with values in \mathbb{R} are the same. It is easy to check that the corresponding symplectic manifold (M, ω) is isomorphic to the symplectic manifold (\mathbb{R}^2, ω_0) , where ω_0 is the standard symplectic form on \mathbb{R}^2 . Unfortunately, for the groups $G \neq \mathrm{SU}(1, 1)$ mentioned above we do not know whether the symplectic manifolds (M, ω) and $(\mathbb{R}^{2n}, \omega_0)$, where $\dim M = 2n$, are isomorphic or not.

The following proposition is known. We do not know a good reference for this; then we give a short proof communicated to us by Yu.A. Neretin.

Proposition 4.3. *For each symmetric Hermitian space $M = G/K$, where G is a connected simple Lie group and K is its maximal compact subgroup, the corresponding Guichardet-Wigner cocycle is non-trivial in the complex $C^*(G, \mathbb{R})$.*

Proof. Let $p : \tilde{G} \rightarrow G$ be the universal cover and let $a = C_{x_0}$ be the Guichardet-Wigner cocycle for the group G . Consider the corresponding to a 2-cocycle \tilde{a} on \tilde{G} induced by p . By construction, the cocycle \tilde{a} is trivial, i.e., there is a smooth function b defined on \tilde{G} such that for any $g, h \in \tilde{G}$ we have $\tilde{a}(g, h) = b(h) - b(gh) + b(g)$.

Assume that the cocycle a is trivial in the complex $C^*(G, \mathbb{R})$, i.e., there exists a function $f : G \rightarrow \mathbb{R}$ such that for $g, h \in G$ we have $a(g, h) = f(h) - f(gh) + f(g)$.

Then the difference $b - f \circ p$ is a homomorphism $\tilde{G} \rightarrow \mathbb{R}$. This homomorphism vanishes near the identity element of \tilde{G} since the group \tilde{G} is simple, and thus it vanishes on the whole of \tilde{G} since \tilde{G} is connected. Then the function f is smooth and the cocycle a is trivial in the complex $C_{\mathrm{diff}}^*(G, \mathbb{R})$. This contradiction proves our statement. \square

5. CASES OF NONTRIVIALITY OF THE COCYCLE C_{x_0} FOR GROUPS OF SYMPLECTOMORPHISMS

Let (M, ω_M) be a non-compact symplectic manifold such that $H^1(M, \mathbb{R}) = 0$ with an exact symplectic form ω_M .

By formula (2), the form ω_M defines a 2-cocycle C_{x_0} for the group $\mathrm{Diff}(M, \omega_M)$ with values in the trivial $\mathrm{Diff}(M, \omega_M)$ -module \mathbb{R} . The aim of this section is to indicate cases when this cocycle is non-trivial and thus the corresponding central extension of the group $\mathrm{Diff}(M, \omega_M)$ by \mathbb{R} is non-split.

Let $M = G/K$ be an Hermitian symmetric space M and let (M, ω) be the corresponding symplectic manifold which we considered in subsection 4.2.

Theorem 5.1. *For the Hermitian symmetric space $M = G/K$ and for the corresponding symplectic manifold (M, ω) the cocycle C_0 on the group $\mathrm{Diff}(M, \omega)$ is non-trivial.*

Proof. Since the group G is a subgroup of the group $\mathrm{Diff}(M, \omega)$ the statement follows from proposition 4.3. \square

Recall that the symplectic manifold (H^2, ω) where ω is the area form is symplectomorphic to (\mathbb{R}^2, ω_0) where ω_0 is the standard symplectic form.

Theorem 5.2. *Let (M, ω) be a non-compact symplectic manifold such that the symplectic form ω_M is exact and let $H^1(M, \mathbb{R}) = 0$. Consider the product $\mathbb{R}^2 \times M$ of the manifold \mathbb{R}^2 and M as a symplectic manifold with the symplectic form $\omega = \omega_0 + \omega_M$. Then for each point $x_0 \in \mathbb{R}^2 \times M$ the cocycle C_{x_0} on the group $\text{Diff}(\mathbb{R}^2 \times M, \omega)$ is non-trivial.*

Proof. Choose $\omega_{M,1} \in \Omega^1(M)$ with $d\omega_{M,1} = \omega_M$ and let $\omega_1 = x dy + \omega_{M,1}$. The group $\text{Diff}(\mathbb{R}^2, \omega_0)$ acting on the first factor \mathbb{R}^2 of $\mathbb{R}^2 \times M$ is naturally included as a subgroup into the group $\text{Diff}(\mathbb{R}^2 \times M, \omega)$. Thus $g^*\omega_1 - \omega_1 = g^*(x dy) - x dy$ for all g in the subgroup $\text{Diff}(\mathbb{R}^2, \omega_0)$. Thus the cocycle C_{x_0} constructed from the form $dx \wedge dy + \omega_M$ on $\mathbb{R}^2 \times M$ restricts to a nontrivial cocycle on the subgroup of $\text{Diff}(\mathbb{R}^2, \omega_0)$ by proposition 4.3 applied to the group $\text{PSL}(2, \mathbb{R})$. \square

We leave to the reader to formulate the corresponding results for other symmetric Hermitian spaces G/K instead of H^2 .

5.3. Problem. Consider an open disk M in the Euclidean plane equipped with the standard area 2-form ω . Is the 2-cocycle C_{x_0} defined by the form ω non-trivial?

6. APPENDIX

In this appendix, for a symplectic manifold (M, ω) we define a 2-cocycle on the Lie algebra $\text{Vect}(M, \omega)$ of locally Hamiltonian or Hamiltonian vector fields, corresponding to the 2-cocycle C_{x_0} on the group $\text{Diff}(M, \omega)$, and study conditions of its nontriviality.

Let \mathfrak{g} be a Lie algebra over \mathbb{R} and let \mathbb{R} be the trivial \mathfrak{g} -module. Denote by $C^p(\mathfrak{g}, \mathbb{R})$ the space of skew-symmetric p -forms on \mathfrak{g} with values in \mathbb{R} . For $c \in C^p(\mathfrak{g}, \mathbb{R})$ and x_1, \dots, x_{p+1} put

$$(4) \quad (\delta^p c)(x_1, \dots, x_{p+1}) = \sum_{i < j} (-1)^{i+j} c([x_i, x_j], x_1, \dots, \hat{x}_i, \dots, \hat{x}_j, \dots, x_{p+1}),$$

where, as usual, \hat{x} means that x is omitted. Then $C^*(\mathfrak{g}, \mathbb{R}) = (C^p(\mathfrak{g}, \mathbb{R}), \delta^p)_{p \geq 0}$ is the complex of standard cochains of the Lie algebra \mathfrak{g} with values in the trivial \mathfrak{g} -module \mathbb{R} and the cohomology $H^*(\mathfrak{g}, \mathbb{R})$ of this complex is the cohomology of the Lie algebra \mathfrak{g} with values in the trivial \mathfrak{g} -module \mathbb{R} .

In particular, there is a bijective correspondence between $H^2(\mathfrak{g}, \mathbb{R})$ and the set of isomorphism classes of central extensions of the Lie algebra \mathfrak{g} by \mathbb{R} .

Let (M, ω) be a symplectic manifold. Denote by $\text{Vect}(M, \omega)$ the Lie algebra of locally Hamiltonian vector fields and by $\text{Vect}_0(M, \omega)$ the Lie algebra of Hamiltonian vector fields on M . For a point $x_0 \in M$ and $X, Y \in \text{Vect}(M, \omega)$ put $c_{x_0}(X, Y) = \omega(X, Y)(x_0)$.

Proposition 6.1. *The function $c_{x_0} : \mathfrak{g}^2 \rightarrow \mathbb{R}$ is a 2-cocycle on the Lie algebra \mathfrak{g} with values in the trivial \mathfrak{g} -module \mathbb{R} . The cohomology class of c_{x_0} is independent of the choice of the point x_0 .*

Proof. The proof is given by direct calculations and is based on the standard formulas $[\mathcal{L}_X, \mathbf{i}_Y] = \mathbf{i}_{[X, Y]}$ and $\mathcal{L}_X = \mathbf{i}_X d + d\mathbf{i}_X$, where \mathbf{i}_X is the operator of the inner product by X and \mathcal{L}_X is the Lie derivative with respect to a vector field X , (see, for example, [6], ch. 4). In particular, we have for any $x \in M$ and $X, Y \in \text{Vect}(M, \omega)$ the following equality

$$(5) \quad c_x(X, Y) - c_{x_0}(X, Y) = - \int_{x_0}^x \mathbf{i}_{[X, Y]} \omega.$$

□

Let G be a Lie group and let \mathfrak{g} be its Lie algebra. We have a natural homomorphism of complexes $C_{\text{diff}}^*(G, \mathbb{R}) \rightarrow C^*(\mathfrak{g}, \mathbb{R})$ (see, for example, [4], ch. 3). In particular, if $c \in C_{\text{diff}}^2(G, \mathbb{R})$, the corresponding cochain $\tilde{c} \in C^2(\mathfrak{g}, \mathbb{R})$ is defined as follows:

$$\tilde{c}(X, Y) = \frac{\partial^2}{\partial t \partial s} (c(\exp tX, \exp sY) - c(\exp sY, \exp tX))_{t=0, s=0}$$

where $X, Y \in \mathfrak{g}$.

Let G be a Lie group of diffeomorphisms of M contained in the group $\text{Diff}(M, \omega)$. Then for the 2-cocycle $c = C_{x_0}$ of section 3, the cocycle \tilde{c} is cohomologous to the restriction of the cocycle c_{x_0} to the Lie algebra \mathfrak{g} of G . Unfortunately, we cannot apply this procedure to the whole group $\text{Diff}(M, \omega)$ and the Lie algebra $\text{Vect}(M, \omega)$. Therefore, the problems of nontriviality of 2-cocycles C_{x_0} on the group $\text{Diff}(M, \omega)$ and c_{x_0} on the Lie algebra $\text{Vect}(M, \omega)$ should be solved independently.

For each $X \in \text{Vect}(M, \omega)$ denote by α_X the closed 1-form such that $\alpha_X = \mathbf{i}_X \omega$. For all vector fields $X, Y \in \text{Vect}(M, \omega)$ we have the following equality:

$$(6) \quad \omega(X, Y) \omega^n = n \alpha_X \wedge \alpha_Y \wedge \omega^{n-1}$$

which can be easily checked in Darboux coordinates.

Denote by X_f a Hamiltonian vector field defined by a function $f \in C^\infty(M)$. Consider the Poisson algebra $\text{P}(M) = \text{P}(M, \omega)$ on (M, ω) , i.e., the algebra $C^\infty(M)$ endowed with the Poisson bracket $\{f, g\} = -\omega(X_f, X_g)$ for $f, g \in C^\infty(M)$.

The map $\text{P}(M) \rightarrow \text{Vect}_0(M, \omega)$ given by $f \rightarrow X_f$ is a homomorphism of Lie algebras which defines an extension of $\text{Vect}_0(M, \omega)$ by \mathbb{R} . It is easy to check that this extension is isomorphic to one given by the cocycle $-c_{x_0}$.

Theorem 6.2. *For a non-compact symplectic manifold (M, ω) the cocycle c_{x_0} on the Lie algebras $\text{Vect}(M, \omega)$ and $\text{Vect}_0(M, \omega)$ is non-trivial.*

Proof. It suffices to prove our statement for the Lie algebra $\text{Vect}_0(M, \omega)$.

First we prove that for each form $\beta \in \Omega^{2n-1}(M)$ there is a unique form $\alpha \in \Omega^1(M)$ such that $\beta = \alpha \wedge \omega^{n-1}$. Indeed, using Darboux coordinates it is easy to check that this has a unique local solution α . These are compatible and we get a global solution by gluing them.

Note that for each form $\alpha \in \Omega^1(M)$ there is a positive integer N and $2N$ functions $f_k, g_k \in C^\infty(M)$ ($k = 1, \dots, N$) such that $\alpha = \sum_{k=1}^N f_k dg_k$ which follows easily from the existence (by dimension theory) of a finite atlas for M .

Since $H^{2n}(M, \mathbb{R}) = 0$ there is a form $\beta \in \Omega^{2n-1}(M)$ such that $\omega^n = d\beta$. Then we have $\omega^n = \sum_{k=1}^N df_k \wedge dg_k \wedge \omega^{n-1}$. By (6) and using this equality we get

$$(7) \quad \sum_{k=1}^N \{f_k, g_k\} = -n.$$

Assume that the extension $P(M) \rightarrow \text{Vect}_0(M, \omega)$ $P(M) \rightarrow \text{Vect}_0(M, \omega)$ is split. Then $P(M)$ is a direct sum of the space of constant functions on M and an ideal isomorphic to $\text{Vect}_0(M, \omega)$ by $P(M) \rightarrow \text{Vect}_0(M, \omega)$. Equality (7) means that these summands have nonzero intersection. This contradiction proves the statement. \square

Now we consider a compact symplectic manifold (M, ω) . It is known that the extension $P(M) \rightarrow \text{Vect}_0(M, \omega)$ is split.

For a closed form α denote by $[\alpha]$ the cohomology class of α . Denote by L the linear map $H^p(M, \mathbb{R}) \rightarrow H^{p+2}(M, \mathbb{R})$ defined by $a \rightarrow a \smile [\omega]$, where $a \in H^p(M, \mathbb{R})$.

Theorem 6.3. *Let (M, ω) be a compact symplectic manifold. The cocycle c_{x_0} on the Lie algebra $\text{Vect}(M, \omega)$ is non-trivial iff the linear map*

$$L^{n-1} : H^1(M, \mathbb{R}) \rightarrow H^{2n-1}(M, \mathbb{R})$$

is not equal zero.

Proof. We may assume that $\int_M \omega^n = 1$. Put for brevity $V = \text{Vect}(M, \omega)$ and $V_0 = \text{Vect}_0(M, \omega)$. Set

$$b(X, Y) = \int_M \omega(X, Y) \omega^n,$$

where $X, Y \in V$. It is easy to check that b is a 2-cocycle on V .

Multiplying both sides of equality (5) by ω^n and integrating over M we get

$$(8) \quad b(X, Y) - c_{x_0}(X, Y) = \int_M \left(\int_{x_0}^X \mathbf{i}_{[X, Y]} \omega \right) \omega^n.$$

Since the right hand side of (8) is a coboundary of a 1-cochain in $C^1(V, \mathbb{R})$, the cocycles c_{x_0} and b are cohomologous. By (6) we have

$$(9) \quad b(X, Y) = n \int_M \alpha_X \wedge \alpha_Y \wedge \omega^{n-1},$$

for any $X, Y \in V$. If $X \in V_0$ the form α_X is exact, and $b(X, Y) = 0$ by (9). This proves (1).

Suppose that the cocycle b is trivial, i.e., there is a linear functional f on V such that for any $X, Y \in V$ we have $b(X, Y) = f([X, Y])$. By [1] we have $[V, V] = [V_0, V_0] = V_0$. This implies $b = 0$. So the cocycle b is trivial iff it equals zero. By (9) and the Poincaré duality this implies that $L^{n-1} = 0$ on $H^1(M, \mathbb{R})$. This proves (2). \square

We know no example when $H^1(M, \mathbb{R}) \neq 0$ and the map $L^{n-1} = 0$. Moreover, if M is a compact Kählerian manifold the map $L^{n-1} : H^1(M, \mathbb{R}) \rightarrow H^{2n-1}(M, \mathbb{R})$ is an isomorphism (see, for example, [11], ch. 4). Thus in this case the cocycle c_{x_0} is non-trivial whenever $H^1(M, \mathbb{R}) \neq 0$.

REFERENCES

- [1] V. I. Arnold, *The one-dimensional cohomology of the Lie algebra of divergence-free vector fields, and the winding numbers of dynamical systems*, Functional. Anal. Appl., **3**, no. 4, (1969), 77-78 (Russian). English translation: Functional. Anal. Appl., **3**, (1969), 319-321.
- [2] J.-L. Dupont, A. Guichardet. *A propos de l'article "Sur la cohomologie réelle des groupes de Lie simples réels"*, Ann. Sci. Ec. Norm. Sup., **11** (1978), 293-296.
- [3] W.T. van Est, *Group cohomology and Lie algebra cohomology in Lie groups. I, II*, Indagationes Mathematicae, **15** (1953), 484-492; 493-504.
- [4] A. Guichardet, *Cohomologie des groupes topologiques et des algèbres de Lie*, Cedric/Nathan, Paris, 1980.
- [5] A. Guichardet, D. Wigner, *Sur la cohomologie réelle des groupes de Lie simples réels*, Ann. Sci. Ec. Norm. Sup., **11** (1978), 277-292.
- [6] C. Godbillon, *Geometrie differentielle et mecanique analytique*, Hermann, Paris, 1969.
- [7] R.S. Ismagilov, *Infinite-dimensional groups and their representations*, in: Proc. Intern. Congress Math. (Warszawa, 1983), 861-875 (Russian).
- [8] S. MacLane, *Homology*, Grundlehren 114, Springer-Verlag, 1963.
- [9] G.D. Mostow, *Cohomology on topological groups and solvmanifolds*, Ann. Math., **73** (1961), 20-48.
- [10] Y.A. Neretin, *Central extensions of groups of symplectomorphisms*, arXiv:math.DG/0406213
- [11] A. Weil, *Introduction à l'étude des variétés kähleriennes*, Hermann, Paris, 1958.

R. S. ISMAGILOV: BAUMAN MOSCOW STATE UNIVERSITY, 2-ND BAUMANSKAYA STR. 5, 107005 MOSCOW, RUSSIA.

E-mail address: ismagil@serv.bmstu.ru

M. LOSIK: SARATOV STATE UNIVERSITY, ASTRAKHANSKAYA 83, 410026 SARATOV, RUSSIA.

E-mail address: LosikMV@info.sgu.ru

P. W. MICHOR: FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, NORDBERGSTRASSE 15, A-1090 WIEN, AUSTRIA; and: ERWIN SCHRÖDINGER INSTITUTE OF MATHEMATICAL PHYSICS, BOLTZMANNGASSE 9, A-1090 WIEN, AUSTRIA.

E-mail address: Peter.Michor@esi.ac.at