

Gluing local gauge conditions in BV quantum field theory

Ezra Getzler

Northwestern University

17 August, 2020

Quantization of the superparticle

The superparticle is a dimensional reduction to $d = 1$ of the Green-Schwarz superstring. It has bosonic fields $x^\mu(t)$ and $p_\mu(t)$, position and momentum in 10-dimensional Minkowski space, and a fermionic field $\theta^a(t)$, a Majorana-Weyl spinor.

The Lagrangian density is

$$L = p_\mu dx^\mu - \frac{1}{2} \eta^{\mu\nu} p_\mu p_\nu dt - \frac{1}{2} \gamma_{ab}^\mu p_\mu \theta^b d\theta^a.$$

The Green-Schwarz superstring is a nonlinear σ -model with target 10-dimensional superspace, with a WZW term associated to the closed three-form with coefficients $\gamma_{\mu ab}$. The “kinetic” term for the fermion needs to be gauge fixed, but this cannot be done in a Lorentz invariant way: on-shell, the bilinear form $\gamma_{ab}^\mu p_\mu$ has rank 8.

The BV formalism allows us to gauge-fix the superparticle by choosing a fixed light-like vector q_μ , and restricting to the submanifold where $c(q)\theta$ and $c(q)\theta^+$ vanish. This is called the **light-cone gauge**.

The BV formalism consists of two parts:

- a solution of the “quantum master equation”, which is obtained from the action of the original classical field theory by coupling to ghosts and extending to higher orders in \hbar by performing renormalization;
- the choice of a Lagrangian in the space of fields and antifields, which may be interpreted as a choice of gauge for the field theory.

We show how to replace the Lagrangian by a “flexible Lagrangian”, a coherent family of Lagrangians defined in different patches and glued together in a manner reminiscent of the proof of the de Rham theorem.

The introduction of allows us to repair a flaw of the light-cone gauge: the gauge-fixed action is degenerate along the ray $p_\mu \propto q_\mu$. This is why we have introduced the idea of a flexible Lagrangian.

All of the results of this talk are taken from

E. G. and Sean W. Pohorence, *Global gauge conditions in the Batalin-Vilkovisky formalism*. To appear. [arXiv:1911.11269](https://arxiv.org/abs/1911.11269)

The Berezinian

Much of differential geometry extends from manifolds to supermanifolds: differentiable maps, Taylor's theorem, the chain rule, the inverse function theorem, the theory of fiber bundles, and the definition of the tangent and cotangent bundles.

We can define the de Rham complex, but there is a big difference: if ξ^a and ξ^b are odd coordinates, the differentials $d\xi^a$ and $d\xi^b$ commute. In particular, the de Rham complex is unbounded. This means that a theory of integration must be based on a different line bundle, due to Berezin.

The definition of this line bundle is based on the geometry of the tangent bundle, which is spanned at a point of U_α by the vector space with basis (x^α even coordinates and ξ^a odd coordinates)

$$\left\{ \frac{\partial}{\partial x^\alpha}, \frac{\partial}{\partial \xi^a} \right\}.$$

A remark on grading

In applications, all supermanifolds in the BV formalism are \mathbb{Z} -graded. In other words, the commutative superalgebras of functions have an auxiliary \mathbb{Z} -grading, which does not in general have anything to do with the parity of the functions. We will ignore this point here.

Lie supergroups

A Lie supergroup is a supermanifold with the structure of a group. The general linear supergroup $GL(n|m)$ is a basic example; it is an open supermanifold of the superspace of all linear maps from $\mathbb{R}^{n|m}$ to itself. We may represent such a matrix in block form

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

with the matrix entries of A and D even, and of B and C odd.

The Berezinian

Just as the tangent bundle of an n -dimensional manifold has structure group $GL(n)$, the tangent bundle of an $n|m$ -dimensional supermanifold has structure group $GL(n|m)$. Given a rational representation of $GL(n|m)$, we may form the associated vector bundle.

The supergroup $GL(n|m)$ has a character

$$\text{Ber} : GL(n|m) \rightarrow GL(1)$$

given by the explicit formula

$$\begin{aligned} \text{Ber} \begin{pmatrix} A & B \\ C & D \end{pmatrix} &= \text{Ber} \begin{pmatrix} I_n & 0 \\ CA^{-1} & I_m \end{pmatrix} \text{Ber} \begin{pmatrix} A & B \\ 0 & D - CA^{-1}B \end{pmatrix} \\ &= \frac{\det(A)}{\det(D - CA^{-1}B)}. \end{aligned}$$

Note that this character really is rational, and not polynomial.

Integration

The density bundle $|\Omega|$ on a supermanifold is the line bundle associated to the character $|\text{Ber}|^{-1}$. Berezin showed that there is a covariantly defined integral

$$\int : \Gamma_c(M, |\Omega|) \rightarrow \mathbb{R}.$$

In local coordinates, a section of $|\Omega|$ has the form

$$f(x, \xi) \left| \frac{dx^1 \dots dx^n}{d\xi^1 \dots d\xi^m} \right|,$$

and its integral is obtained by integrating the function

$$\frac{\partial^m f(x, 0)}{\partial \xi^1 \dots \partial \xi^m}$$

with respect to x .

Odd symplectic supermanifolds

A two-form ω determines a bundle map from TM to T^*M ; it is **non-degenerate** if this map is invertible, and **symplectic** if it is non-degenerate and closed. The symplectic form is **odd** if its total parity is odd.

Darboux's theorem holds, with essentially the same proof as usual: there are coordinates $(x^1, \dots, x^n, \xi^1, \dots, \xi^n)$ such that

$$\omega = \sum_{a=1}^n dx^a \wedge d\xi^a.$$

In particular, the dimension of an odd symplectic supermanifold is $n|n$.

The odd cotangent bundle $M = \Pi T^*X$, where X is a supermanifold, is an odd symplectic supermanifold. In physics, the coordinates on the base are called **fields**, and the coordinates along the fibres are called **antifields**.

The structure group of the tangent bundle of an odd symplectic supermanifold does not have a conventional notation: Manin calls it $\Pi\text{Sp}(n|n)$.

The following observation was made by Khvadaverdian and Voronov (though it is implicit in the work of Batalin and Vilkovisky).

Lemma

There is a character $\text{Ber}^{1/2} : \Pi\text{Sp}(n|n) \rightarrow \text{GL}(1)$ whose square is the restriction of the Berezinian along $\Pi\text{Sp}(n|n) \subset \text{GL}(n|n)$.

In fact, we have the formula

$$\text{Ber}^{1/2} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(A).$$

The line bundle associated to the character $|\text{Ber}|^{-1/2}$ is the half-density bundle $|\Omega|^{1/2}$. Its nontriviality is a measure of the failure of Liouville's theorem for odd symplectic supermanifolds. (By contrast, the symplectic group $\text{Sp}(2n)$ is a subgroup of the special linear group $\text{SL}(2n)$.)

Khudaverdian's differential operator

In terms of a Darboux coordinate system, we may write a half-density as

$$f(x, \xi) |dx^1 \dots dx^n|.$$

There is an operator defined on sections $\Gamma(M, |\Omega|^{1/2})$, given in a Darboux coordinate system by the formula

$$\Delta f = \sum_{a=1}^n \frac{\partial^2 f(x, \xi)}{\partial x^a \partial \xi^a} |dx^1 \dots dx^n|.$$

It is clear that $\Delta^2 = 0$ (since $\partial^2 f / \partial \xi^a \partial \xi^b + \partial^2 f / \partial \xi^b \partial \xi^a = 0$).

Theorem (Khudaverdian)

The operator Δ is independent of the Darboux coordinate system.

The quantum master equation

A nowhere-vanishing section $s \in \Gamma(M, |\Omega|^{1/2})$ of the half-density bundle satisfying $\Delta s = 0$ is called an **orientation** of M . (This terminology is due to Behrend and Fantechi.) A supermanifold with a pair (ω, s) of an odd symplectic form and an orientation may be called a **Batalin–Vilkovisky supermanifold**. This encapsulates the basic structures of quantum field theory.

Suppose that in a Darboux coordinate system, the orientation s is given by the formula $e^S |dx^1 \dots dx^n|$ for some function S , called the **action**. Then the equation $\Delta s = 0$ becomes

$$\sum_{a=1}^n \left(\frac{\partial^2 S}{\partial x^a \partial \xi^a} + \frac{\partial S}{\partial x^a} \frac{\partial S}{\partial \xi^a} \right) = 0.$$

This is the **quantum master equation** of Batalin and Vilkovisky.

In quantum field theory, we may expand S in powers of \hbar (the [loop expansion](#)):

$$S = \sum_{g=0}^{\infty} \hbar^{g-1} S_g.$$

The quantum master equation becomes

$$\sum_{a=1}^n \left(\frac{\partial^2 S_{g-1}}{\partial x^a \partial \xi^a} + \sum_{h=0}^g \frac{\partial S_h}{\partial x^a} \frac{\partial S_{g-h}}{\partial \xi^a} \right) = 0.$$

The leading term of this equation is the [classical master equation](#)

$$\sum_{a=1}^n \frac{\partial S_0}{\partial x^a} \frac{\partial S_0}{\partial \xi^a} = 0.$$

This equation turns out to encapsulate all generalized gauge symmetries of classical actions. . . .

Integrals on Batalin-Vilkovisky manifolds

In modern language, the problem that Batalin and Vilkovisky answer is: how do we construct linear forms $Z : \Gamma_c(M, |\Omega|^{1/2}) \rightarrow \mathbb{R}$ such that

$$Z(\Delta s) = 0$$

for all half-densities s ? Such a linear form is called a **trace**; they are used by Batalin and Vilkovisky to define the functional integrals of quantum field theories.

Kontsevich and Schwarz interpret traces as generalized sections $Z \in \Gamma^{-\infty}(M, |\Omega|^{1/2})$ satisfying $\Delta Z = 0$.

A **Lagrangian submanifold** of an odd symplectic manifold M is a supermanifold $\iota : L \hookrightarrow M$ properly embedded in M such that the symplectic form vanishes on restriction to L , and induces an isomorphism between the tangent and normal bundles of L .

It follows that the half-density bundle $|\Omega_M|_M^{1/2}$ restricts along L to the density bundle $|\Omega_L|$.

Batalin and Vilkovisky's main result is that for any $\sigma \in \Gamma_c(M, |\Omega|^{1/2})$,

$$\int_L \iota^* \Delta \sigma = 0.$$

That is, integration of half-forms on M along a Lagrangian L defines a trace.

Families of Lagrangians

In the remainder of this talk, we will give a more general construction of traces on the complex $\Gamma_c(M, |\Omega|^{1/2})$ of half-densities. The key ingredient is a formula of Mikhailov and Schwarz for the variation of \int_{L_t} when L_t varies in a family of Lagrangians.

The tangent space at L to the space of Lagrangians in a symplectic manifold equals the space of closed 1-forms on L . The same result holds for Lagrangians in odd symplectic manifolds.

There is one small difference: the resulting 1-form has odd parity, and the odd de Rham cohomology vanishes, hence the 1-form is actually exact, and represented by an odd function on L . Furthermore, this function is uniquely determined.

If L_t is a one-parameter family of Lagrangians, let η_t be the corresponding family of functions. The 1-form $\eta_t dt$ has even total parity, since the 1-form dt has odd parity in the de Rham complex of the interval.

Let $\Delta^k = \{0 \leq t_1 \leq \dots \leq t_k \leq 1\} \subset \mathbb{R}^k$ be the k -dimensional simplex.

If $\iota : L \times \Delta^k \rightarrow M$ is an k -dimensional family of Lagrangians, we may apply the above construction to obtain a 1-form η on $L \times \Delta^k$, of even total parity.

Let δ be the de Rham differential along Δ^k .

Theorem

If $\sigma \in \Gamma_c(M, |\Omega|^{1/2}) \otimes \Omega^*(\Delta^k)$, we have

$$\delta \int_L e^{-\eta} \iota^* \sigma = \int_L e^{-\eta} \iota^* (\delta + \Delta) \sigma.$$

The proof uses the Darboux–Weinstein theorem for Lagrangians in odd symplectic manifolds; this is proved in the same way as the usual Darboux–Weinstein theorem, using Moser’s trick. The theorem of Mikhailov and Schwarz is an infinitesimal version of this formula.

Flexible Lagrangian submanifolds

Let M be an odd symplectic supermanifold, and let $\mathcal{U} = \{U_\alpha\}$ be a locally finite open cover, with partition of unity $\{\phi_\alpha\}$.

By a **flexible Lagrangian**, we mean a collection of parametrized families of proper embeddings

$$\begin{array}{ccc} L_{\alpha_0 \dots \alpha_k} \times \Delta^k & \xrightarrow{\quad} & U_{\alpha_0 \dots \alpha_k} \times \Delta^k \\ & \searrow \quad \swarrow & \\ & \Delta^k & \end{array}$$

one for each multi-index $\alpha_0 \dots \alpha_k$.

These families are required to glue together appropriately as we approach the faces. For example, as $t \rightarrow 0$, $\iota_{\alpha_0 \alpha_1}(t)$ approaches the intersection of $\iota_{\alpha_0} : L_{\alpha_0} \rightarrow M_{\alpha_0}$ with U_{α_1} , while as $t \rightarrow 1$, it approaches the intersection of $\iota_{\alpha_1} : L_{\alpha_1} \rightarrow M_{\alpha_1}$ with U_{α_0} .

The Thom–Whitney complex

We borrow an idea from Sullivan's rational homotopy theory. The Thom-Whitney complex $\text{Tot } |\Omega|^{1/2}(\mathcal{U})$ is the subspace of

$$\prod_{k=0}^{\infty} \prod_{\alpha_0 \dots \alpha_k} \Gamma(U_{\alpha_0 \dots \alpha_k}, |\Omega|^{1/2}) \otimes \Omega^*(\Delta^k)$$

compatible with all of the simplicial maps. (This is formally similar to the definition of the geometric realization of a simplicial space: this similarity is no accident.)

The differential on this complex is $\delta + \Delta$, and the complex is quasi-isomorphic to the Čech complex of the sheaf $|\Omega|^{1/2}$, and hence, since these sheaves are fine, to the $\mathbb{Z}/2$ -graded complex $\Gamma(M, |\Omega|^{1/2})$ with differential Δ .

The main theorem

Let $\Phi_{\alpha_0 \dots \alpha_k}$ be the differential operator on $|\Omega|^{1/2}(U_{\alpha_0 \dots \alpha_k})$ given by the formula

$$\Phi_{\alpha_0 \dots \alpha_k} = \frac{1}{k+1} \sum_{i=0}^k [\Delta, \phi_{\alpha_0}] \dots [\Delta, \phi_{\alpha_{i-1}}] \phi_{\alpha_i} [\Delta, \phi_{\alpha_{i+1}}] \dots [\Delta, \phi_{\alpha_k}].$$

Theorem

Let σ_{\bullet} be a compactly supported element of the Thom–Whitney complex $\text{Tot } |\Omega|^{1/2}(\mathcal{U})$. Define the linear form

$$Z(\sigma_{\bullet}) = \sum_{k=0}^{\infty} (-1)^k \sum_{\alpha_0 \dots \alpha_k} \int_{\Delta^k} \int_{L_{\alpha_0 \dots \alpha_k}} e^{-\eta_{\alpha_0 \dots \alpha_k}} l_{\alpha_0 \dots \alpha_k}^* (\Phi_{\alpha_0 \dots \alpha_k} \sigma_{\alpha_0 \dots \alpha_k}).$$

Then Z is a trace: $Z((\delta + \Delta)\sigma_{\bullet}) = 0$.

The **rank** of a Batalin-Vilkovisky quantization is the degree of the Lagrangian density L as a function of the **antifields**. Rank one theories, in which the dependence on antifields is linear, reduce to the BRS formalism.

In the special case of rank one theories, our main result is already presented in the article

C. Becchi, G. Giusto and C. Imbimbo, *The functional measure of gauge theories in the presence of Gribov horizons*, [arXiv:hep-th/9811018](https://arxiv.org/abs/hep-th/9811018)

The superparticle is a rank two theory, in which the action has a quadratic dependence on the antifields. In this case, our results appear to be new.