

Programme on
“Measured Group Theory”
January 18 - March 18, 2016

organized by

**Miklos Abert (Hungarian Academy of Sciences, Budapest), Goulmara Arzhantseva (U Vienna),
Damien Gaboriau (ENS Lyon), Thomas Schick (U Göttingen), Andreas Thom (TU Dresden)**

MINI-COURSE
“On the Graham Higman group”
by Prof. Lev Glebsky (U Autónoma de San Luis Potosí, Mexico)

• **Wednesday, January 20, 2016**

10:00 – 12:00

• **Friday, January 22, 2016**

10:00 – 12:00

• **Wednesday, January 27, 2016**

10:00 – 12:00

• **Friday, January 29, 2016**

10:00 – 12:00

Abstract

Let $H_{m,k} = \langle a_0, \dots, a_{m-1} \mid \{a_i^{-1} a_{i-1} a_i = a_{i-1}^k, i = 0, \dots, m-1\} \rangle$, here $i-1$ is taken mod m . In 1951 Graham Higman introduced the group $H_{4,2}$ as an example of a finitely presented infinite group without finite quotients. Recently, some researchers are speculating if the group $H_{4,2}$ could serve as an example of non-sofic group. During the lectures we study some properties of $H_{m,k}$ trying to answer how plausible these speculations are.

Topics to consider

- The groups $B(1, k) = \langle a, b \mid a^b = a^k \rangle \sim \mathbb{Z} \ltimes \mathbb{Q}_k$ and $\langle a_0, a_1, a_2 \mid a_1^{a_2} = a_1^k, a_2^{a_3} = a_2^k \rangle$ as ‘building blocks’ for $H_{m,k}$. Their properties.
- The G. Higman proof (in a little bit more general context) that $H_{4,2}$ has no finite quotients.
- $H_{m,k}$, $k > 2$, $m > 4$ is not residually finite but nevertheless has a lot of finite p -quotients for $p \mid (k-1)$. Construction of pro- p ‘extension’ of $H_{m,k}$.
- $B(1, k)$ as a subgroup of $H_{m,k}$. Exponent-like modular functions.

All talks take place at the ESI, Boltzmann Lecture Hall!